

New Role for DCR-1/Dicer in *Caenorhabditis elegans* Innate Immunity against the Highly Virulent Bacterium *Bacillus thuringiensis* DB27

Igor latsenko, Amit Sinha, Christian Rödelsperger, Ralf J. Sommer

Max Planck Institute for Developmental Biology, Department of Evolutionary Biology, Tübingen, Germany

Bacillus thuringiensis produces toxins that target invertebrates, including *Caenorhabditis elegans*. Virulence of *Bacillus* strains is often highly specific, such that *B. thuringiensis* strain DB27 is highly pathogenic to *C. elegans* but shows no virulence for another model nematode, *Pristionchus pacificus*. To uncover the underlying mechanisms of the differential responses of the two nematodes to *B. thuringiensis* DB27 and to reveal the *C. elegans* defense mechanisms against this pathogen, we conducted a genetic screen for *C. elegans* mutants resistant to *B. thuringiensis* DB27. Here, we describe a *B. thuringiensis* DB27-resistant *C. elegans* mutant that is identical to *nasp-1*, which encodes the *C. elegans* homolog of the nuclear-autoantigenic-sperm protein. Gene expression analysis indicated a substantial overlap between the genes downregulated in the *nasp-1* mutant and targets of *C. elegans dcr-1*/Dicer, suggesting that *dcr-1* is repressed in *nasp-1* mutants, which was confirmed by quantitative PCR. Consistent with this, the *nasp-1* mutant exhibits RNA interference (RNAi) deficiency and reduced longevity similar to those of a *dcr-1* mutant. Building on these surprising findings, we further explored a potential role for *dcr-1* in *C. elegans* innate immunity. We show that *dcr-1* mutant alleles deficient in microRNA (miRNA) processing, but not those deficient only in RNAi, are resistant to *B. thuringiensis* DB27. Furthermore, *dcr-1* overexpression rescues the *nasp-1* mutant's resistance, suggesting that repression of *dcr-1* determines the *nasp-1* mutant's resistance. Additionally, we identified the collagen-encoding gene *col-92* as one of the downstream effectors of *nasp-1* that play an important role in resistance to DB27. Taken together, these results uncover a previously unknown role for DCR-1/Dicer in *C. elegans* antibacterial immunity that is largely associated with miRNA processing.

Genetically tractable model organisms, including the nematode *Caenorhabditis elegans*, have provided detailed insights into the origin and fundamental principles of immunity. More than a decade of research on *C. elegans* has yielded a wealth of knowledge about its innate immune response to various pathogens, uncovering multiple signaling pathways critical to *C. elegans* survival after exposure to pathogens (for reviews, see references 1 to 7). Once activated, these pathways induce the expression of an array of antimicrobial effectors that differ, depending on the pathogen present (8–11). A variety of microbes, including bacteria (12), fungi (13), viruses (14), and microsporidia (15), have been used in *C. elegans* infection studies, with the main focus on etiological agents of human diseases. In some cases, work on *C. elegans* has led to the identification of conserved virulence factors of human pathogens (16, 17).

Among the natural pathogens of *C. elegans*, *Bacillus thuringiensis* has been extensively studied and *B. thuringiensis* likely coexists and coevolves with its host in the natural environment (18). *B. thuringiensis* produces a variety of pore-forming toxins (PFTs), called Cry toxins, that target the intestinal cells of insects (19) and nematodes (20). The ability of some Cry toxins to target *C. elegans* has been exploited to use this nematode as an *in vivo* system for studying PFTs (21), and several *C. elegans* defense mechanisms against PFTs have been discovered (22–27).

Although the *C. elegans* model has provided invaluable insight into the immune defense against pathogens, a comparative approach with other nematodes would enhance our understanding of the host response to pathogens. One nematode that has been used extensively for comparative studies with *C. elegans* is *Pristionchus pacificus*. In contrast to *C. elegans*, which is often found on compost heaps and rotten fruit (28), *P. pacificus* is found in tight association with scarab beetles (29). Not surprisingly, therefore, these two nematodes show different resistance patterns

(30-32) and transcriptional responses to pathogens (11). For example, C. elegans dies when fed the human opportunistic bacteria Pseudomonas aeruginosa and Staphylococcus aureus and B. thuringiensis Cry5B toxin, whereas P. pacificus is resistant (20, 30). Moreover, we have previously shown (31) that out of 768 naturally isolated Bacillus strains, only 20 were pathogenic and show distinct patterns of virulence for C. elegans and P. pacificus. The most extreme difference was seen with regard to B. thuringiensis strain DB27. DB27 kills C. elegans in less than 24 h, representing one of the fastest killers of C. elegans known to date, whereas P. pacificus is resistant to this bacterial strain (31). Our previous study (31) also indicated that the determinants of B. thuringiensis DB27 virulence for C. elegans and also the mechanisms of C. elegans resistance to this pathogen are unknown. Specifically, we have shown that the bre and daf-2 mutants, which are known to be resistant to B. thuringiensis PFT (22) and other bacterial pathogens (33), respectively, are as susceptible to *B. thuringiensis* DB27 as wild-type worms are (31). These findings suggest that potentially novel mechanisms are required to provide defense against this highly virulent Bacillus strain.

Here, we report a genetic screen for B. thuringiensis DB27-

Received 3 June 2013 Returned for modification 10 July 2013 Accepted 1 August 2013 Published ahead of print 5 August 2013 Editor: J. F. Urban, Jr. Address correspondence to Ralf J. Sommer, ralf.sommer@tuebingen.mpg.de. Supplemental material for this article may be found at http://dx.doi.org/10.1128 /IAI.00700-13. Copyright © 2013, American Society for Microbiology. All Rights Reserved. doi:10.1128/IAI.00700-13 resistant mutants of C. elegans and show that a gene represented by two alleles is identical to the nasp-1 gene C09H10.6. Further characterization revealed that the nasp-1 mutant shows a reduced life span and reduced fecundity and is RNA interference (RNAi) deficient. Microarray analysis uncovered an unexpected enrichment of dcr-1-regulated genes in nasp-1 mutants. We explored a potential role for *dcr-1* in *C. elegans* innate immunity and show that *dcr-1* allele mutants deficient in microRNA (miRNA) processing, but not those that are only deficient in RNAi, are resistant to B. thuringiensis DB27. Transgenic expression of dcr-1 in nasp-1 mutants can rescue the resistance phenotype of nasp-1 mutants. We identified the collagen-encoding gene col-92 as one of the downstream effectors of *nasp-1* and *dcr-1*. These findings describe NASP-1 as a novel C. elegans innate immunity regulator and identify a previously unknown role for DCR-1 in the C. elegans antibacterial immune response.

MATERIALS AND METHODS

Nematodes and bacterial strains. *C. elegans* strains were maintained on nematode growth medium (NGM) agar plates with *Escherichia coli* OP50 as a food source. The following strains were kindly provided by the *Caenorhabditis* Genetics Center (University of Minnesota): wild-type Bristol (N2), Hawaiian mapping strain (CB4856), DA453 *eat-2(ad453)*, WM49 *rde-4(ne301)*, WM27 *rde-1(ne219)*, CB189 *unc-32(e189)*, BB1 *dcr-1(ok247)/unc-32(e189) III*, YY11 *dcr-1(mg375)*, VC1138 *drsh-1(ok369)*, VC446 *alg-1(gk214)*, BA1 *fer-1(hc1)*, and BA15 *fer-15(hc15)*. *btr-1/ C09H10.9(tu439)* and *btr-1(tu440)* mutant worms were generated in this study and backcrossed to wild-type worms four times before analysis. Double mutants were generated by using standard procedures (34). *Serratia marcescens* C2 was isolated from Reunion Island, *S. aureus* Newman and *P. aeruginosa* PA14 were purchased from the Deutsche Sammlung von Mikroorganismen und Zellkulturen, and *B. thuringiensis* DB27 was isolated from dung beetles (31).

Genetic screen for *B. thuringiensis* DB27-resistant nematodes. After ethyl methanesulfonate (EMS) mutagenesis (50 mM for 4 h at 20°C), single parental (P_0) mothers were separated onto NGM OP50 plates and allowed to lay 10 to 30 eggs. F_1 mothers were allowed to lay F_2 progeny that, upon reaching the L4 stage, were placed onto plates with pathogenic *B. thuringiensis* DB27. After 24 h of incubation at 25°C, single worms that survived pathogen exposure were transferred to NGM OP50 plates and allowed to reproduce. These candidates were retested several times in survival assay to confirm the resistance phenotype.

C. elegans killing assays. Each bacterium was grown overnight in a shaking incubator at 30°C in Luria-Bertani (LB) broth, except *S. aureus* and *P. aeruginosa*, which were grown at 37°C. A 100- μ l volume of the culture was spread to the edges of a 6-cm NGM plate to prevent worms from escaping and incubated overnight. Three to six independent replicates of 20 adult worms per plate were exposed to pathogens and monitored for survival. Survival assays were repeated multiple times and conducted at 25°C. Nematodes were transferred once a day to fresh plates and considered dead when they failed to respond to touch. We note that the exact nature of the killing assay (inoculum size etc.) might have an influence on the quantitative killing results reported in this study.

Complementation test, mapping, sequencing, and rescue of *tu439.* Complementation tests were performed by testing F_1 progeny from the crosses between *btr-2(tu438)* mutant males and *dpy-5;tu439* or *dpy-5; tu440* mutant hermaphrodites. To test if *tu439* and *tu440* are allelic, F_1 progeny from a cross between *tu439* mutant males and *dpy-5;tu440* mutant hermaphrodites were tested. For mapping, the *btr-1* mutant was crossed to Hawaiian strain CB4856 and approximately 400 F_2 cross progeny were isolated and screened for *B. thuringiensis* DB27 resistance. DNA was isolated from a total of 96 *tu439*-positive recombinant progeny and used for restriction fragment length polymorphism (RFLP)–single-nucleotide polymorphism (SNP) mapping (35). SNP data from the most informative recombinants identified a 160-kb region on chromosome II between the markers F38A3 and F37H8. Whole-genome sequencing of *btr-1* mutants revealed that both *btr-1* alleles contain two identical mutations in the mapping interval, which were found in the gene C09H10.6. For transformation rescue, a 2.3-kb PCR fragment, covering upstream and downstream regions of the C09H10.6 transcript, was amplified from wild-type genomic DNA and injected together with a *sur-5::gfp* plasmid as a coinjection marker into a *btr-1* mutant as described elsewhere (36).

For dcr-1, dpy-5, and col-92 overexpression, respective transcripts with presumptive regulatory regions were PCR amplified from wild-type genomic DNA, purified, and used for injection together with the sur-5::gfp plasmid as a coinjection marker. For intestine-specific expression, a 3.5-kb cpr-1 promoter fragment was amplified from wild-type DNA and fused with the dcr-1 transcript by using Phusion High-Fidelity DNA polymerase (NEB) according to the manufacturer's instructions. The resulting construct was used for btr-1 microinjection. Reporter gene constructs for gene expression analysis were created by the PCR fusion method as described previously (37). A nasp-1::GFP (green fluorescent protein) translational reporter was obtained by PCR amplification of the nasp-1 transcript with the presumptive promoter region from wild-type genomic DNA, which in the next step was PCR fused to GFP, which was amplified from the pPD95.75 plasmid. A col-92 transcriptional reporter was generated in the same way, with the exception that only a 1.4-kb col-92 promoter region was fused to GFP. The resulting constructs were purified and used for injection together with the pRF4 plasmid (rol-6). Several independent lines were obtained and tested for all of the transgenic experiments described here.

Mutant identification by whole-genome sequencing. We sequenced a genomic library of the btr-1(tu439) mutant on the Illumina Genome Analyzer II platform and obtained 7,398,278 read pairs $(2 \times 100 \text{ bp})$. Raw reads were aligned with the C. elegans genome assembly (WS235) by using bwa (version 0.5.9-r16) (38). After the removal of duplicated reads to exclude false variant calls due to PCR amplification biases, 98.8% of the C. elegans assembly was covered by at least one read, with a genome-wide mean coverage of 15×. Variants were called by using SAMtools (version 0.1.18-r982:295) (39). In total, we identified 2,092 homozygous base substitutions with a SAMtools quality score of at least 20, of which six variants were located within the 160-kb mapping interval on chromosome II. Filtering for variants with a potential deleterious effect (nonsynonymous substitution, premature stop codons, and mutations near splice sites) could exclude three variants as intergenic and intronic, respectively. Two of the remaining three nonsynonymous substitutions were found to affect the nasp-1 gene (amino acid changes, L53S and D307N), and one substitution at genomic position II:11069603, G to A (WS235), was found to cause an S-to-F amino acid exchange in various alternative transcripts of unc-53. A rescue experiment confirmed that mutations responsible for the phenotype occurred in the nasp-1 gene.

Pumping rate measurement. Pharyngeal pumping was counted for 1 min starting on the first day of adulthood. Worms were placed on *B. thuringiensis* DB27 plates at 25°C and left undisturbed for 3 h before measurement. All worms remained on food during the period of observation. The *eat-2(ad453)* mutant, known to have low pumping, was used as a control.

Brood size measurement. Brood size was determined by picking L4 worms (one per plate) and transferring them to fresh plates every 24 h until egg laying ceased. The offspring were counted 2 days after the mothers were removed from the plates. Ten worms were used per strain per experiment, and experiments were repeated twice.

Life span assay. Life span assays were performed on NGM plates containing 0.1-mg/ml 5'-fluorodeoxyuridine (FUDR) to prevent progeny from hatching at 20°C. Plates were seeded with concentrated OP50 and allowed to dry overnight. A synchronized population of L4 worms grown on OP50 was placed onto the plates and scored every 48 h. For *col-92* RNAi life span assays, NGM plates containing 50 µg/ml carbenicillin, 1 mM isopropyl-β-D-thiogalactoside (IPTG), and 0.1 mg/ml FUDR were seeded with *E. coli* HT115 expressing *col-92* double-stranded RNA (dsRNA) (experiment) or *E. coli* HT115 harboring the empty vector (control). A synchronized population of L4 worms grown on *col-92* RNAi bacteria or vector-containing bacteria were placed onto the RNAi plates with FUDR and scored every 48 h. Upon depletion of food, worms were transferred to fresh plates. Worms were considered dead when no response to touch was observed. Approximately 100 worms were used per strain per experiment, and experiments were repeated twice.

RNAi. To generate specific gene knockdowns, we used RNAi by feeding nematodes with E. coli expressing dsRNA that is homologous to a target gene. E. coli harboring the empty vector was used as a control. E. coli strain HT115 harboring the appropriate vector was grown in LB broth containing 100 µg/ml ampicillin at 37°C overnight. Bacteria were plated onto NGM plates containing 50 µg/ml carbenicillin and 1 mM IPTG and allowed to grow overnight at 37°C. Three adult hermaphrodites were placed on each RNAi plate and allowed to lay eggs. These eggs were grown on RNAi plates for 3 days at 22°C to the adult stage before being transferred to plates with B. thuringiensis DB27 for pathogenicity assay. All of the RNAi clones used in this study are part of the RNAi library obtained from Source BioScience UK Limited (Nottingham, United Kingdom). cdc-25 RNAi plates were prepared as described above. However, worms were grown at 25°C to the adult stage. cdc-25.1 encodes a CDC25 phosphatase homolog that affects embryonic viability and is necessary for cell proliferation in the germ line. In brief, gravid worms were laid on cdc-25.1 RNAi plates for 4 h and then transferred to similar plates for an additional 4 h of egg laying. After that, gravid hermaphrodites were removed and eggs were left to hatch and grow in the presence of cdc-25.1 RNAi to produce sterile worms. cdc-25 RNAi by injection was performed as described below. For mom-2 RNAi, three adult hermaphrodites were placed on mom-2 RNAi plates and allowed to lay eggs. These eggs were grown on RNAi plates at 20°C to the adult stage. Single adults were subsequently placed onto freshly prepared mom-2 RNAi plates and allowed to lay eggs for 24 h and then removed from the plates. After 16 h, the total amounts of eggs and hatched larvae were scored and viability was expressed as follows: (number of hatched larvae/total number of eggs) \times 100. The mom-2 coinjection experiments were conducted at 20°C as described previously (40).

dsRNA for injection experiments was prepared by *in vitro* transcription. DNA templates were PCR amplified from plasmids corresponding to the RNAi clones with T7 primers or PCR amplified from *C. elegans* genomic DNA with gene-specific primers with attached T7 promoters in case the gene is not present in the available RNAi library. PCR products were used for dsRNA synthesis in a single reaction with the TranscriptAid T7 high-yield transcription kit (Thermo Scientific). Concentrations of dsRNAs were determined with a spectrophotometer, and the quality and size of dsRNAs were assessed by gel electrophoresis. Young hermaphrodites were injected in the gonads with 300 to 600 ng/µl of dsRNA of the candidate gene. Given that *dpy-5* RNAi produces a clear morphological phenotype but has no effect on survival, *dpy-5* dsRNA was used as a positive control for injection and as a negative control in survival assays. Adult F₁ worms were used in the pathogen survival assay.

RNA collection for microarray experiments. Synchronized populations of wild-type and *tu439* mutant *C. elegans* worms were obtained by hypochlorite treatment and allowed to grow to the young adult stage on *E. coli* OP50 at 20°C. Worms were exposed to *B. thuringiensis* DB27 for 4 h and then picked into TRIzol reagent for RNA extraction. Four biological replicates were collected for each experimental condition and the control condition. For each biological replicate, about 200 young adult hermaphrodites were picked into 1 ml of TRIzol (Invitrogen) and total RNA was extracted according to the manufacturer's instructions and purified further by phenol-chloroform-isoamyl alcohol precipitation. The RNA pellet was suspended in RNase-free water and assessed on a NanoDrop spectrophotometer for quantity and RNA quality.

Microarray experiments and data analysis. Oligonucleotide microarrays for *C. elegans* containing ~43,000 unique probes for ~20,000 *C*. *elegans* genes were obtained from Agilent Technologies (NCBI GEO accession number GPL10094). All experiments are in a two-color format, where Cy5 and Cy3 dye-labeled cRNAs from experimental and control samples are cohybridized on the same microarray. The four biological replicates per experiment included two dye swap experiments to account for differences in dye labeling. Experimental procedures and data analysis were essentially the same as described before (41). Microarrays corresponding to three of the four biological replicates passed the quality criteria and were used in the final analysis. Genes with a false-discovery rate (FDR)-corrected *P* value of ≤ 0.05 and an absolute fold change greater than or equal to a cutoff of 1.5 on a \log_2 scale were called differentially expressed.

Expression cluster enrichment analysis. The list of microarray expression profiles in which a given *C. elegans* gene is known to be differentially expressed can be extracted from the Expression Cluster section of the WormBase (42) gene summary page for each gene. We retrieved all of the available expression clusters for *C. elegans* genes from the WormBase website (www.wormbase.org) and used them to identify expression clusters enriched in our expression profile, an approach that we have successfully used previously to analyze genome-wide trends in expression profiles (11, 41). *P* values for expression cluster enrichment in each expression profile were computed with a two-by-two Fisher exact test. An FDR-corrected *P* value cutoff of 0.05 was used as the significance threshold.

qRT-PCR. Synchronized *C. elegans* worms at the L4 stage were exposed to *B. thuringiensis* DB27 for 4 h before harvesting. Nematodes were collected by washing plates with M9 buffer, and RNA was extracted with TRIzol reagent. cDNA was synthesized with the SuperScript III kit (Invitrogen). Quantitative real-time PCR (qRT-PCR) was conducted with the LightCycler 480 SYBR green I master kit (Roche) on a LightCycler 480 real-time PCR instrument (Roche) in a 96-well format. Relative fold changes in transcripts were calculated by the comparative cycle threshold (C_T) method after normalization to *snb-1*, which has been used previously in qRT-PCR studies of *C. elegans* innate immunity (43–45). C_Ts were determined by the Roche LightCycler software. All samples were run in triplicate. The sequences of the primers used are available upon request.

Statistical analysis. Kaplan-Meier nonparametric comparisons and a log rank test were used for statistical analysis of *C. elegans* survival in the presence of pathogens and for life span analysis. In each case, curves represent the combined data from at least three independent experiments. Statistical analysis of the difference between two values was performed with Student's *t* test. Statistical significance was set at $P \le 0.05$.

Microarray data accession number. Raw and processed data from all of the experiments described here are available under accession number GSE43905 in the NCBI Gene Expression Omnibus database (http://www .ncbi.nlm.nih.gov/geo/).

RESULTS

Forward genetic screen for B. thuringiensis DB27-resistant (btr) mutants. B. thuringiensis DB27 was isolated from a dung beetle (31) and was selected for detailed investigation because of its high virulence for C. elegans. This strain is 100% lethal in just 16 h under standard assay conditions (Fig. 1A). Our previous studies indicated that all of the bre and daf-2 mutants tested were as susceptible to *B. thuringiensis* DB27 as wild-type worms are (31). In contrast to the observed susceptibility of C. elegans, the second model nematode organism, P. pacificus, is fully resistant to B. thuringiensis DB27 (31). To uncover a potential resistance mechanism and to better understand the specificity of B. thuringiensis virulence for certain nematode species, we conducted a forward genetic screen for mutants of *C. elegans* with strong resistance to *B*. thuringiensis DB27. Specifically, we screened for mutants that would extend C. elegans survival on monoxenic cultures of B. thuringiensis DB27 by at least a factor of 3, in order to select for major effectors. From 1,800,000 EMS-mutagenized gametes, we isolated



FIG 1 Differences in the survival of wild-type and *btr* mutant worms upon exposure to different pathogenic bacteria. (A) *C. elegans* wild-type (wt) N2 is highly susceptible to *B. thuringiensis* (BT) DB27, in contrast to standard *E. coli* OP50, which was used as a control (P < 0.0001). (B) Survival of *btr* mutants on monoxenic cultures of *B. thuringiensis* DB27. All alleles showed significantly (P < 0.0001) enhanced survival compared to that of the wild type. (C) *tu439* mutants show increased resistance to *S. aureus* compared to that of the wild type (P < 0.0001). (D, E) *tu439* mutants are as susceptible as wild-type worms to *P. aeruginosa* (D, P > 0.05) and to *S. marcescens* (E, P > 0.05). For survival curves, the numbers of live versus dead worms were scored over time. The number of worms alive (N nematodes) is plotted as a function of time. The data shown are means \pm the standard errors of the means. For each condition in the survival assay, 60 to 80 worms (20 per plate) were used for each of multiple independent biological replicates.

three *btr* (<u>B</u>. *thuringiensis*-resistant) mutants that exhibited strong resistance to <u>B</u>. *thuringiensis*-mediated killing (Fig. 1B). Complementation tests revealed that these mutants fall into two complementation groups (data not shown) with two alleles (tu439 and tu440) of a locus provisionally named btr-1 and a single allele (tu438) of a second locus called btr-2. Here we further characterize btr-1.

btr-1(tu439) mutants are resistant to additional pathogens and have a reduced life span. Considering that several pathogenresistant *C. elegans* mutants are resistant to multiple pathogens (33), we exposed *btr-1(tu439)* mutant worms to additional pathogens to further characterize the specificity of the mutant. We found that *tu439* mutants exhibited stronger resistance to the Gram-positive pathogen *S. aureus* than wild-type worms did (Fig. 1C). In contrast, *tu439* mutants showed wild-type survival on the Gram-negative pathogens *P. aeruginosa* and *S. marcescens* (Fig. 1D and E). These results indicate that *tu439* confers resistance to two, but not all, of the pathogens tested.

To explore additional functions, we tested several physiologi-

cal parameters of tu439 mutant worms. First, we estimated the pumping rate as an indirect measurement of bacterial uptake to elucidate bacterial consumption. The pumping rate of the tu439 mutant on DB27 is similar to that of the wild type, whereas the control eat-2(ad453) mutant shows the expected reduction of pumping in our assay system (Fig. 2A). Thus, pharyngeal pumping is unaffected, suggesting that pathogen consumption is not reduced in mutant worms. To rule out the possibility that bacteria might persist in the intestines of wildtype but not mutant worms, we fed worms with DB27 for 4 h, which is sufficient to observe accumulation of bacteria in the gut (see Fig. S1A in the supplemental material). We then shifted the worms to OP50 for 2 h and measured the DB27 CFU. Neither the wild-type nor the mutant worms showed a difference in their DB27 CFU counts after the shift to OP50, suggesting that the persistence of DB27 is not extended in wildtype worms (see Fig. S1B). Second, we observed that tu439 mutants exhibit reduced fertility (see Fig. S1C). A similar phenotype was also annotated for other pathogen-resistant mu-



FIG 2 *tu439* mutants exhibit increased resistance to DB27 independently of reduced fertility and longevity. (A) The pumping rate of *tu439* mutants shows no difference from that of the wild type (P > 0.05). The control *eat-2* mutant strain shows the expected lower pumping rate than that of the wild type (P < 0.0001). At least 10 worms were scored per strain. The data shown are means ± the standard errors of the means. (B) *fer-1* and *fer-15* mutants show no difference (P > 0.05) from wild-type worms on a lawn of *B. thuringiensis* (BT) DB27. The data shown are means ± the standard errors of the means. For each condition in survival assays, 60 to 80 worms (20 per plate) were used for each of multiple independent biological replicates. (C) Wild-type and *tu439* nematodes were injected with *cdc-25* interfering dsRNA to obtain sterile worms or with control interfering dsRNA (see Materials and Methods) and exposed to DB27. *cdc-25* RNA is ginificantly higher (P < 0.0001) for *tu439*; cdc-25 mutant worms is significantly higher (P < 0.0001) for *tu439*; mutants is significantly higher (P < 0.0001) lower than that of *tu439* mutants is *significantly* higher (P < 0.0001) lower than that of *tu439* mutants is significantly of *tu439*; mutants is significantly (P < 0.001) lower than that of wild-type worms on standard food (*E. coli* OP50). At least 100 worms were used per strain per experiment, and each experiment was done twice. The data shown are means ± the standard errors of the means.

tants (46–48) and suggests that there is a tradeoff between reproduction and an elevated immune response. This raises the possibility that tu439 confers increased resistance to *B. thuringiensis* DB27 because of diminished fecundity.

To elucidate this further, we assessed the survival of sterile mutants exposed to *B. thuringiensis* DB27. First, we investigated fecundity in combination with matricidal internal hatching of eggs by comparing the survival of wild-type worms to that of *fer-1* and *fer-15* mutants (47). *fer* mutants have a germ line but do not suffer from matricidal effects due to the lack of fertilization since their sperm production is affected (49). Both *fer* mutants are as susceptible to *B. thuringiensis* DB27 as wild-type worms are, suggesting that matricidal hatching of eggs does not contribute to *B. thuringiensis* DB27 killing and that increased resistance of *tu439* to DB27 is not due to a lack of matricide (Fig. 2B). This finding is further supported by the fact that males are susceptible to DB27 (not shown).

In a second set of experiments, we used *cdc-25* RNAi, which also causes sterility. *cdc-25* RNAi led to enhanced survival of wild-type worms exposed to *B. thuringiensis* DB27 (Fig. 2C). However, *cdc-25* RNAi in a *tu439* background resulted in a significant increase in resistance to DB27 that was stronger than the resistance of single *tu439* mutant or wild-type *cdc-25* RNAi worms (Fig. 2C). The cumulative effect of the two mutations suggests that the underlying molecular mechanisms are independent and that *tu439*-mediated resistance to *B. thuringiensis* DB27 is not due simply to

reduced fecundity. Additionally, in contrast to sterile mutants, the *tu439* mutant is not resistant to *S. marcescens* and *P. aeruginosa* (Fig. 1D and E). Finally, we measured the life span of *tu439* mutants, given that life span extension often correlates with increased pathogen resistance in *C. elegans* (33, 47). Interestingly, we found that *tu439* mutants fed on *E. coli* OP50 show a slight but significant life span reduction in comparison to wild-type worms (Fig. 2D). These results show a novel relationship between life span and immunity and suggest that *tu439* mutant worm resistance to bacterial infection is not a simple consequence of extended longevity. Taken together, *tu439* mutant worms are resistant to several Gram-positive bacteria, have a reduced life span and fecundity, and show that immunity can be mechanistically uncoupled from aging.

btr-1 is identical to the *C. elegans nasp-1* gene. To clone *tu439*, we used RFLP-SNP mapping (35) and next-generation sequencing (50). In brief, we mapped *btr-1* to an approximately 160-kb region on chromosome II between the markers F38A3 and F37H8. Whole-genome sequencing of both alleles revealed mutations in the C09H10.6 gene, which has been described as the *C. elegans* homolog of the gene for human nuclear-autoantigenic-sperm protein (*nasp-1*) (Fig. 3A). *nasp-1* has previously been implicated in *C. elegans* female development (51).

Interestingly, the two alleles appear to contain two identical mutations, which might be due to the fact that they were derived from a single mother. The first mutation is a T-to-C substitution



FIG 3 Identification of the *tu439* allele. (A, top) Exon-intron structure of the C09H10.9 gene, adapted from WormBase (WS235), with the positions of mutations indicated by arrows. Numbers indicate positions on chromosome II where single-nucleotide mutations occurred. (A, bottom) Schematic domain structure of the C09H10.9 protein. The black box shows the location of the TPR domain, which is not affected in either of the *btr-1* mutants. The dashed lines highlight the amino acid sequences flanking the mutation sites. The black arrows show the position and type of amino acid substitution. (B) Rescue experiment. Transgenic expression of *nasp-1* under the control of its endogenous promoter restores wild-type (wt) susceptibility to *B. thuringiensis* (BT) DB27 to the *tu439* mutant. (C) *tu439* mutant resistance to *S. aureus* is rescued by transgenic expression of *nasp-1* under the control of its endogenous promoter restores of *nasp-1* under the control of *B. thuringiensis* DB27 of wild-type wersus the transgenic line). The data shown are means ± the standard errors of the means. (D) The survival on a lawn of *B. thuringiensis* DB27 of wild-type worms grown on dsRNA for C09H10.6, which corresponds to *tu439*, is significantly (P < 0.001) greater than that of worms grown on dsRNA as a vector control. (E) The data shown are means ± the standard errors of the errors of the means. For each condition in survival assays, 60 to 80 worms (20 per plate) were used for each of multiple independent biological replicates.

in the second exon of *C. elegans nasp-1* resulting in a leucine-toserine substitution at amino acid position 53, whereas the second mutation (a G-to-A change in exon 3) leads to a substitution from aspartic acid to asparagine at amino acid position 307 (Fig. 3A). To confirm that the resistance phenotype of *tu439* is due to these mutations in C09H10.6, we performed rescue and RNAi experiments. First, transgenic expression of a wild-type copy of C09H10.6 under the control of its endogenous promoter in the *tu439* mutant background resulted in almost complete restoration of wild-type susceptibility to *B. thuringiensis* DB27 (Fig. 3B) and *S. aureus* (Fig. 3C). Additionally, transgenic expression of C09H10.6 partially rescued the diminished reproduction of the *tu439* mutant (not shown). Second, C09H10.6 RNAi increases nematode resistance to *B. thuringiensis* DB27 (Fig. 3D) and *S. aureus* (Fig. 3E) in a way similar to that of *tu439*, providing additional evidence that the mutant is allelic to C09H10.6. Also, we noticed that *nasp-1* RNAi worms have slower growth and less fecundity at 25°C than vector-treated worms. The high conservation of the amino acids that were mutated (see Fig. S2 in the supplemental material), especially D307, suggests that our *nasp-1* mutant is a strong reduction-of-function allele. Also, the similarity between the *nasp-1* RNAi and the mutant phenotype suggests that the isolated alleles represent reduction-of-function rather than neomorphic mutations. In summary, we conclude that *btr-1(tu439)* is identical to the *nasp-1*-like gene C09H10.6. Following standard *C. elegans* no-menclature rules, we rename *btr-1* as *nasp-1*.



FIG 4 nasp-1 mutant exhibits RNAi deficiency and requires PMK-1 but not DAF-16 for DB27 resistance. (A) pmk-1 knockdown upon RNAi by injection leads to hypersusceptibility in wild-type (wt) worms (P < 0.0001 for wt; dpy-5 versus wt; pmk-1 worms) and abolishes the resistance of the nasp-1 mutant (P < 0.0001for nasp-1;dpy-5 versus nasp-1;pmk-1 mutant worms) to B. thuringiensis (BT) DB27 below the level of wild-type survival (P < 0.05 for nasp-1;pmk-1 versus wt worms). daf-16 knockdown upon RNAi by injection has no effect on nasp-1 mutant (P > 0.05 for nasp-1; dpy-5 versus nasp-1; daf-16 mutant worms) resistance to B. thuringiensis DB27. vhp-1 RNAi significantly (P < 0.0001) increases the survival of wild-type worms on a lawn of B. thuringiensis DB27 compared to control dpy-5 RNAi, which was used as a positive control for injection and as a negative control in a survival assay (see Materials and Methods). (B) pmk-1 knockdown upon RNAi by feeding has no effect on nasp-1-mediated resistance to B. thuringiensis DB27 (P > 0.05), while pmk-1 knockdown upon RNAi by injection abolishes the resistance of the *nasp-1* mutant (P < 0.0001). The data shown are means \pm the standard errors of the means. For each condition in survival assays, 60 to 80 worms (20 per plate) were used for each of multiple independent biological replicates. (C) nasp-1 mutant is RNAi deficient. Wild-type and nasp-1 and rde-4 (positive control) mutant worms and nasp-1 rescuing line worms were fed E. coli expressing dsRNA of mom-2, which is essential for viability. Viability was assessed as the proportion of hatched larvae relative to the total number of eggs laid. Both rde-4 and nasp-1 mutants show significantly (P < 0.0001) higher survival of progeny than the wild type, indicating that both mutants are resistant to RNAi-mediated mom-2 lethality. nasp-1 overexpression restores the *nasp-1(tu439)* mutant's susceptibility to *mom-2*-mediated lethality. The data shown are means \pm the standard errors of the means. (D) Wild-type hermaphrodites were coinjected with each candidate dsRNA and mom-2 dsRNA. rde-1 RNAi was used as a positive control, while unc-32 RNAi was used as a negative control. *nasp-1* RNAi significantly increases F_1 viability (P < 0.0001) relative to that of the negative control or *mom-2* RNAi alone. The data shown are means \pm the standard errors of the means.

The p38 MAPK pathway is required for the nasp-1 mutant's resistance. Next, we searched for potential interactions of nasp-1 with known pathogen response regulators to further explore its biological role. First, we tested the p38 mitogen-activated protein kinase (MAPK) pathway (52) by performing injection of pmk-1 dsRNA into a nasp-1 mutant background. After pmk-1 depletion, the resistance of the nasp-1 mutant to B. thuringiensis DB27 is completely abolished, even below the wild-type level (Fig. 4A). Similarly, overactivation of PMK-1 by vph-1 RNAi (53) strongly increased the survival of C. elegans on B. thuringiensis DB27 (Fig. 4A). In contrast, injection of dpy-5 as a control dsRNA had no effect on the survival of wild-type and *nasp-1* mutant worms, although it did result in a dpy phenotype (Fig. 4A). Together, these results suggest that the effects of pmk-1 and vhp-1 on survival are gene specific and that *nasp-1* potentially may function in parallel with the p38 MAPK pathway. In contrast, RNAi of daf-16, another C. elegans innate immunity regulator (33), had no effect on the nasp-1 mutant's resistance to B. thuringiensis DB27 (Fig. 4A), indicating that nasp-1 does not confer resistance to B. thuringiensis DB27 by acting through daf-16/FOXO. This result also provides an additional argument that the *nasp-1* mutant's resistance is independent of reproduction, because increased pathogen resistance of most of the sterile mutants requires *daf-16* (47).

nasp-1 mutants are deficient for RNAi by feeding. When performing the RNAi experiments described above, we noticed that a knockdown of genes by RNAi in a nasp-1 mutant background was only possible by injection of dsRNA, suggesting that the nasp-1 mutant exhibits a sid phenotype (54). In contrast, RNAi by feeding in the nasp-1 mutant background did not have the same effect (Fig. 4B). Specifically, *pmk-1* RNAi by feeding did not change the nasp-1 mutant's resistance phenotype, whereas pmk-1 RNAi by injection abolished the nasp-1 mutant's resistance to B. thuringiensis DB27 completely (Fig. 4B). Similarly, we observed that dpy-5 RNAi by injection resulted in a dpy phenotype in a nasp-1 mutant background, whereas dpy-5 RNAi by feeding did not (data not shown). These observations provide first evidence that nasp-1 mutants might be defective in their RNAi response; however, they do not exhibit a complete rde phenotype but rather show similarity to sid mutants.

To further explore this phenotype, we fed *nasp-1* mutants with *E. coli* expressing dsRNA against *mom-2*, a gene essential for viability (40, 55). Almost no viable progeny were obtained from wild-

type worms, whereas nearly 80% of the nasp-1 mutant progeny survived mom-2 RNAi by feeding (Fig. 4C). Transgenic expression of nasp-1 under the control of its endogenous promoter in the tu439 mutant background resulted in almost complete restoration of the wild-type response to mom-2-mediated lethality. rde-4 mutants, which are known to be RNAi deficient, were used as a control and showed nearly 100% viability (Fig. 4C). Additionally, we coinjected wild-type worms with dsRNA of mom-2 and/or dsRNA of *nasp-1* or control dsRNA and scored the viability of the progeny. The survival of progeny indicates that RNAi against nasp-1 enables worms to overcome the lethality of mom-2 RNAi (Fig. 4D). Worms coinjected with dsRNA of a positive-control gene, rde-1, showed 97% survival, indicating strong protection from mom-2 lethality by rde-1 RNAi. Coinjection with dsRNA of unc-32 or other genes dispensable for RNAi resulted in a very low survival rate. In contrast, coinjection of nasp-1 dsRNA significantly rescued the lethality caused by mom-2 dsRNA. Thus, in contrast to RNAi mutants, such as *rde-1* RNAi mutants, that are resistant to RNAi independently of the delivery method (56), nasp-1 mutants are still able to perform RNAi when dsRNA is injected into the germ line. In this, they are similar to sid mutants (54), suggesting that they function primarily in the uptake or spreading of dsRNA rather than in modulation of the core RNAi machinery.

To provide further insight into this phenotype, we generated transgenic worms expressing a *nasp-1*-encoded translational protein fusion with GFP to study the site of action of *nasp-1*. Interestingly, expression was localized primarily in the pharynx (see Fig. S3 in the supplemental material). Expression was seen in the metacorpus and terminal bulb of the pharynx with a clear intracellular localization. Given that the pharynx is exposed to pathogens, this observation further supports the idea that *nasp-1* functions in innate immunity; however, the link to the sid phenotype remains unknown. Taken together, these results support the observation that *nasp-1* is required for a robust RNAi response.

The transcriptional response of nasp-1 mutants to B. thuringiensis DB27 reveals a strong overlap with dcr-1 mutants. To identify genes differentially regulated in nasp-1 mutants when they are exposed to B. thuringiensis DB27, we used whole-genome microarrays on an Agilent platform. Microarray analysis revealed that differential expression in nasp-1 mutants mostly resulted in downregulation (247 genes), whereas only 31 genes were more strongly expressed in *nasp-1* mutants than in wild-type worms (see Data Set S1 in the supplemental material, FDR-corrected P value of ≤ 0.05 , fold change cutoff of 1.5 on a log₂ scale). We validated these results by selecting a total of 12 down- and upregulated genes and confirmed their expression by qRT-PCR (Fig. 5A). Interestingly, 58 of the 247 downregulated genes were previously identified as genes that are induced by B. thuringiensis DB27, in comparison to OP50, in wild-type worms (11) (Fig. 5B). These results, together with the finding that most expression changes in nasp-1 mutants result in transcriptional repression, suggest that NASP-1 normally acts as an activator of transcription.

Surprisingly, further characterization of downregulated genes in *nasp-1* mutants revealed a highly significant overlap with genes regulated by different components of the RNAi machinery (57), such as *rde-4*, *rde-3*, and *rrf-1* (see Fig. S4 in the supplemental material). The most significant enrichment was found for genes previously annotated as targets of the only *C. elegans* dicer gene, *dcr-1*. Specifically, 116 of the 247 *nasp-1*-repressed genes were previously identified as *dcr-1*-regulated genes (Fig. 5C). This strong overlap is further supported by quantitative PCR experiments showing that *dcr-1* expression is itself reduced in *nasp-1* mutants compared to that in the wild type (Fig. 5A). These findings are consistent with the reduced life span and RNAi deficiency of *nasp-1* mutants (see above). Thus, an intriguing possibility is that repression of DCR-1 function is responsible for the increased pathogen resistance observed in *nasp-1* mutants.

dcr-1 is involved in innate immunity through its role in miRNA processing. The unexpected correlation between downregulated genes in nasp-1 and dcr-1 mutants encouraged us to look for further interactions between these two loci. We hypothesized that if reduced DCR-1 function in nasp-1 mutants were responsible for their increased resistance to *B. thuringiensis* DB27, dcr-1 mutants should also be more resistant to the pathogen. Indeed, when we tested several dcr-1 alleles, we found that the mutant for presumptive null allele dcr-1(ok247) showed greater survival on *B. thuringiensis* DB27 than the wild type (Fig. 6A). This is unlikely to be due to altered reproduction because the dcr-1(ok247) mutant is not resistant to S. marcescens (see Fig. S5A in the supplemental material) and P. aeruginosa (see Fig. S5B) and exhibits a reduced life span (58), in contrast to sterile mutants (59, 60). As the dcr-1(ok247) deletion allele was marked with a recessive allele of *unc-32* to facilitate the isolation of *dcr-1(ok247)* homozygous worms (57) and considering that the unc-32 mutant was shown to have increased resistance to Erwinia carotovora and Photorhabdus luminescens (8), we also tested the survival of unc-32 mutants on B. thuringiensis DB27 and found that they have a wildtype response to the pathogen (Fig. 6A). Furthermore, transgenic expression of *dcr-1* in a *dcr-1(ok247)* mutant restored wild-type susceptibility to B. thuringiensis DB27 (Fig. 6A). Together, these findings indicate a novel role for *dcr-1* in innate immunity to the pathogen B. thuringiensis DB27.

Interestingly, the effect of *dcr-1* on pathogen resistance is allele specific. We found that survival on B. thuringiensis DB27 was unchanged in the mutant with the activating dcr-1(mg375) allele (Fig. 6A). The dcr-1(mg375) allele has an impaired helicase domain and small interfering RNA (siRNA) production and, consequently, enhanced RNAi, whereas miRNA synthesis is intact (61). To further test the idea that resistance to *B. thuringiensis* DB27 is associated with altered miRNA processing, we investigated the effects of the *rde-4* and *rde-1* mutations on pathogen survival. rde-1 and rde-4 play a role in RNAi but not in miRNA processing (57), and they were shown to produce a reduced life span but normal survival on S. marcescens, although the immune response is upregulated (57). Indeed, *rde-1* and *rde-4* mutants did not show increased resistance to B. thuringiensis DB27 (Fig. 6B). In contrast, mutants with changes in the drsh-1 and alg-1 genes, which are involved in miRNA processing (62, 63), but not in RNAi, exhibited increased resistance to B. thuringiensis DB27 (Fig. 6C). To prove that the resistance of *drsh-1* and *alg-1* mutants is not due to impaired germ line development and fertility, we rendered both mutants sterile via cdc-25 RNAi and compared their survival to that of control vector-treated worms. The survival of alg-1 mutants was significantly increased after cdc-25 RNAi compared to that of vector-treated *alg-1* mutants and compared to wild-type worms treated with either the empty vector or *cdc-25* RNAi (Fig. 6D), indicating that alg-1 and cdc-25 mutants exhibit resistance via distinct mechanisms and that *alg-1*-mediated resistance to *B*. thuringiensis DB27 is not simply a consequence of altered fertility. However, we did not observe any differences in survival between



FIG 5 Transcriptional response of *nasp-1* mutant worms to *B. thuringiensis* DB27 infection. (A) qRT-PCR confirmation of microarray results. Wild-type (wt) and *nasp-1* mutant worms were exposed to *B. thuringiensis* DB27 for 4 h and used for RNA isolation. The data shown are the average of two independent RNA isolations. Each transcript was measured in triplicate and normalized to a control gene. Error bars show the standard errors of the means. (B) Venn diagram of the genes downregulated in the *nasp-1* mutant and upregulated in wild-type worms upon exposure to *B. thuringiensis* DB27 (11). The overlap is significant (P = 1.11E-15). (C) Venn diagram showing the overlap between genes downregulated in the *nasp-1* mutant (47). The overlap is significant (P = 3.70E-79).

drsh-1 vector- and *drsh-1 cdc-25* RNAi-treated worms (Fig. 6E), suggesting that the increased resistance of *drsh-1* mutants to *B. thuringiensis* DB27 is indeed a secondary effect of sterility. Taken together, these results suggest that the impairment of DCR-1 function in miRNA processing confers resistance to *B. thuringiensis* DB27. Furthermore, these data, together with the downregulation of *dcr-1* expression in *nasp-1* mutants, support the conclusion that the resistance of *nasp-1* mutants is due to reduced DCR-1 activity.

nasp-1 acts through DCR-1/Dicer. The experiments described above are consistent with the hypothesis that *dcr-1* is genetically downstream of *nasp-1*. This conclusion is further supported by previous expression profiling of *dcr-1* mutants, which did not show any change in the expression of *nasp-1* in *dcr-1* mutants (57). To further test this hypothesis, we investigated whether overexpression of *dcr-1* would abolish the increased resistance of the *nasp-1* mutant to *B. thuringiensis* DB27. Indeed, *dcr-1* overexpression under the control of its endogenous promoter significantly reduced the survival of *nasp-1* mutant worms on *B. thuringiensis* DB27 (Fig. 6F). Additionally, *dcr-1* overexpression restored the reduced life span of *nasp-1* mutants (see Fig. S6 in the supplemental material). Given that *C. elegans dcr-1* is ex-

pressed at high levels in the intestine (58), which is also the major site of infection by ingested pathogens, we investigated whether upregulation of *dcr-1* specifically in the intestine could affect the *nasp-1* mutant's pathogen resistance. When we overexpressed *dcr-1* under the control of the intestine-specific *cpr-1* promoter, we noticed a significant decrease in the *nasp-1* mutant's survival on *B. thuringiensis* DB27 (Fig. 6F). Thus, the previously unknown role of DCR-1 in *C. elegans* innate immunity is genetically downstream of *nasp-1* and acts through the *C. elegans* intestine.

NASP-1 and Dicer resistance to *S. aureus* is due to impaired fecundity. Considering that the *nasp-1* mutant shows increased resistance to *S. aureus*, we investigated whether it is also mediated by the same mechanism that is employed in the defense against *B. thuringiensis* DB27. Given that the *nasp-1* mutant has reduced fecundity, a common feature of *S. aureus*-resistant mutants (47), we first tested if the *nasp-1* mutant's resistance to *S. aureus* depends on reproduction. As shown in Fig. 7A, *nasp-1* mutants injected with *cdc-25* interfering dsRNA survived *S. aureus* exposure longer than control-injected worms. However, their survival was similar to that of wild-type worms injected with *cdc-25* interfering dsRNA (Fig. 7A). Thus, *nasp-1* mutants do not have an additive effect with *cdc-25* RNAi for *S. aureus* resistance, as is the case for *B.*



FIG 6 Repression of DCR-1 activity in miRNA processing but not in RNAi results in increased resistance to *B. thuringiensis* DB27. (A) The survival of different *dcr-1* alleles on a lawn of *B. thuringiensis* (BT) DB27 in comparison with wild-type (wt) and *nasp-1* mutant worms. *dcr-1(ok247)* mutant worms are significantly (P < 0.0001) more resistant to *B. thuringiensis* DB27 than wild-type worms are. *dcr-1(mg375)* mutant worms show wild-type susceptibility to the pathogen (P > 0.05). Transgenic expression of *dcr-1* rescues the increased resistance of the *dcr-1(ok247)* mutant to *B. thuringiensis* DB27. (B) RNAi-defective *rde-4(ne301)* and *rde-1(ne219)* mutants show survival comparable to that of wild-type worms (P > 0.05) upon exposure to *B. thuringiensis* DB27. (C) Survival of miRNA-deficient mutants upon exposure to *B. thuringiensis* DB27. *alg-1(gk214)* and *drsh-1(ok369)* mutants show resistance to DB27 greater than that of wild-type worms (P < 0.0001). (D, E) Wild-type, *alg-1* mutant (D), and *drsh-1* mutant (E) nematodes were fed with *cdc-25* RNAi vector to obtain sterile worms or with empty RNAi vector and exposed to DB27. *cdc-25* RNAi significantly increases the resistance of wild-type (P < 0.0001 for wt;vector versus wt;*cdc-25* worms) and *alg-1* (D) (P = 0.002 for *alg-1*;vector versus *alg-1;cdc-25* worms but not *drsh-1* mutants (E) (P > 0.05 for *drsh-1*;vector versus *drsh-1;cdc-25* worms). The survival of *alg-1;cdc-25* mutant worms is significantly different (P < 0.05) for that of wt;*cdc-25* worms. The survival of *alg-1;cdc-25* worms of *dcr-1* abolishes the resistance of the *nasp-1* mutant to *B. thuringiensis* DB27. The resistance of *nasp-1* mutants (P < 0.0001) for each condition in survival assays, 60 to 80 worms (20 per plate) were used for each of multiple independent biological replicates. The data shown are means \pm the standard errors of the means.

thuringiensis DB27 (Fig. 2C). These findings suggest that *nasp-1* and *cdc-25* RNAi increase resistance to *S. aureus* via the same mechanism and the *nasp-1* mutant's resistance to *S. aureus* is very likely due to partial sterility. In contrast, *nasp-1* and *cdc-25* RNAi function in independent pathways for *B. thuringiensis* DB27 resistance (Fig. 2).

This difference between *S. aureus* and *B. thuringiensis* DB27 was further explored by studying the allele specificity of Dicer mutants. There was no allele specificity in terms of Dicer resistance to *S. aureus* since both *dcr-1(ok247)* and *dcr-1(mg375)* mutant worms were more resistant to *S. aureus* than wild-type worms were (Fig. 7B). Also, *dcr-1* mutant resistance can be rescued by

transgenic overexpression of dcr-1 under the control of its endogenous promoter (Fig. 7B), indicating that Dicer might play a role in the *C. elegans* defense against *S. aureus*. Considering the reproductive defects of dcr-1 mutants, which may contribute to increased resistance to *S. aureus*, we used cdc-25 RNAi to sterilize dcr-1(mg375) mutant worms. The survival of mg375 mutant worms was greater than that of vector-treated worms but was not significantly different from that of wild-type worms treated with cdc-25 RNAi, indicating the absence of a cumulative effect of dcr-1and cdc-25 (Fig. 7C). Thus, the dcr-1 mutant's resistance to *S. aureus* is due to diminished reproduction, as in the case of the *nasp-1* mutant. We also assessed the survival of RNAi- and



FIG 7 nasp-1- and dcr-1-mediated resistance to S. aureus is due to partial sterility. (A) Wild-type (wt) and nasp-1 mutant nematodes were injected with cdc-25 interfering dsRNA to obtain sterile worms or with control dsRNA (see Materials and Methods) and exposed to S. aureus. cdc-25 RNAi significantly increases the resistance of wild-type (P < 0.0001 for wt;vector versus wt;cdc-25 worms) and nasp-1 mutant (P < 0.0001 for nasp-1;control versus nasp-1;cdc-25 worms) worms. The survival of *nasp-1;cdc-25* mutant worms is not significantly different (P > 0.05) from wt;cdc-25 worms. The data shown are means \pm the standard errors of the means. (B) Both dcr-1 mutants ok247 and mg375 grown at 25°C to induce sterility exhibit increased resistance to S. aureus (P < 0.0001 for ok247 mutant versus wt worms; P < 0.001 for mg375 mutant versus wt worms), which can be rescued by transgenic expression of dcr-1. The ok247 and mg375 mutants do not differ in the level of resistance (P > 0.05). The data shown are means \pm the standard errors of the means. (C) Wild-type and dcr-1(mg375) mutant nematodes were fed with cdc-25 RNAi vector to obtain sterile worms or with empty RNAi vector (see Materials and Methods) and exposed to S. aureus. cdc-25 RNAi significantly increases the resistance of wild-type (P < 0.0001 for wt;vector versus wt;cdc-25 worms) and mg375 (P = 0.018 for mg375;vector versus mg375;cdc-25 mutant worms) worms. The survival of mg375; cdc-25 mutant worms is not significantly different (P > 0.05) from that of wt; cdc-25 worms. The data shown are means ± the standard errors of the means. (D) Survival of wild-type worms and RNAi- and miRNA-deficient mutants upon exposure to S. aureus. Only drsh-1(ok369) mutant worms showed greater S. aureus resistance than wild-type worms (P < 0.0001). The data shown are means \pm the standard errors of the means. (E) Wild-type and drsh-1 mutant nematodes were fed with cdc-25 RNAi vector to obtain sterile worms or with empty RNAi vector and exposed to S. aureus. cdc-25 RNAi significantly increases the resistance of wild-type (P < 0.0001 wt;vector versus wt;cdc-25 worms) but not drsh-1 mutant (P > 0.05drsh-1; vector versus drsh-1; cdc-25 mutant worms) worms. The survival of drsh-1; cdc-25 mutant worms is not significantly different (P > 0.05) from that of wt;cdc-25 worms. For each condition in survival assays, 60 to 80 worms (20 per plate) were used for each of multiple independent biological replicates. The data shown are means \pm the standard errors of the means.

miRNA-deficient mutants on *S. aureus*. Both *rde-1* and *rde-4* mutants showed a wild-type response to *S. aureus* (Fig. 7D). The *alg-1* mutant, which has increased survival on *B. thuringiensis* DB27, exhibited survival similar to that of the wild type on *S. aureus* (Fig. 7D), while only the *drsh-1* mutant was more resistant to *S. aureus* infection (Fig. 7D). However, as in the case of *B. thuringiensis* DB27, the increased resistance seen is the consequence of sterility since *cdc-25* RNAi did not further increase the survival of *drsh-1* mutants (Fig. 7E). Taken together, these results show that the

nasp-1 mutant's resistance to *S. aureus* is very likely due to impaired fecundity and at this stage it is impossible to disconnect immune and reproductive mechanisms that confer the *nasp-1* mutant's resistance to *S. aureus*.

The collagen gene *col-92* acts downstream of *nasp-1* and confers resistance to DB27. Finally, we wanted to uncover the molecular mechanisms of *nasp-1*- and *dcr-1*-mediated resistance to *B*. *thuringiensis* DB27. For this, we hypothesized that the repression of *dcr-1* activity in *nasp-1* mutants leads to overactivation of effec-



FIG 8 Collagen gene *col-92* acts downstream of *nasp-1* and is required for resistance to *B. thuringiensis* (BT) DB27. (A) Some of the DB27 response transcripts (selected from reference 11) show high expression in *dcr-1* mutant worms as determined by qRT-PCR. The data shown are averages of two independent RNA isolations. Each transcript was measured in triplicate and normalized to a control gene. Error bars show the standard errors of the means. (B) Injection of *col-92* interfering dsRNA significantly (P < 0.001) suppresses the *nasp-1* mutant's resistance to *B. thuringiensis* DB27 compared to the injection of a control gene (*dpy-5*). The data shown are means \pm the standard errors of the means. (C) Injection of *col-92* interfering dsRNA significantly (P = 0.021) decreases the *nasp-1* mutant's resistance to *S. aureus* compared to the injection of a control gene (*dpy-5*). The data shown are means \pm the standard errors of the means. (D) Injection of *col-92* increases the standard errors of the means. (D) RNAi knockdown of *col-92* increases the susceptibility of wild-type (wt) worms to *B. thuringiensis* DB27 infection (P < 0.01). The data shown are means \pm the standard errors of *dpy-5* significantly (P < 0.02) increases the survival of worms exposed to *B. thuringiensis* DB27. The data shown are means \pm the standard errors of the means. (F) Mechanism by which *nasp-1* may regulate the innate immune response to *B. thuringiensis* DB27.

tor genes, which make *C. elegans* more resistant to infection. Consistent with this idea, we found that several *B. thuringiensis* DB27responsive genes are more strongly expressed in *nasp-1* and *dcr-1* mutants than in the wild type (Fig. 8A). To test this hypothesis further, we used RNAi to knock down *nasp-1*-upregulated genes individually (see Data Set S1 in the supplemental material) in the *nasp-1* mutant background. While the knockdown of the majority of these genes resulted in no obvious effect on *nasp-1*-mediated resistance (see Fig. S7 in the supplemental material), *col-92* RNAi produced significant suppression of *nasp-1*-mediated resistance to DB27 (Fig. 8B; see Fig. S7) and *S. aureus* (Fig. 8C). Furthermore, knockdown of *col-92* in wild-type worms makes them hypersusceptible to *B. thuringiensis* DB27 (Fig. 8D), *S. aureus*, and *P. aeruginosa* (see Fig. S8A and B) but not to *S. marcescens* (see Fig. S8C). However, the life span of *col-92* RNAi worms was not affected (see Fig. S8D), suggesting that its hypersusceptibility to bacterial infection is not a result of impaired fitness. In contrast, *col-92* overexpression under the control of its endogenous promoter increases resistance to *B. thuringiensis* DB27 (Fig. 8E) and *S. aureus* (see Fig. S8E) infections. At the same time, overexpression of *dpy-5*, used as a control, resulted in no obvious effects on resistance to *B. thuringiensis* DB27 (Fig. 8E) and *S. aureus* (see Fig. S8E). Moreover, *dpy-5* RNAi and *col-93* RNAi had no effect on *C. elegans* survival upon exposure to *B. thuringiensis* DB27, indicating that the effect of *col-92* is specific and is unlikely to be due only to simple alteration of cuticle structure.

To shed more light on the immune function of *col-92*, we studied the expression of the gene. It should be noted that COL-92 is a

member of a nematode-specific family of collagens, characterized by a cuticle collagen N-terminal domain. Transgenic lines harboring a *col-92*–GFP transcriptional fusion construct showed that the gene is expressed predominantly in the hypodermis (see Fig. S9 in the supplemental material). Therefore, although *col-92* is a downstream target of NASP-1, its immune function awaits further characterization and might be associated with cuticle. Thus, the previously uncharacterized collagen gene *col-92* plays a pivotal role in the *C. elegans* defense against bacterial infection and represents a downstream target of NASP-1 and DCR-1.

DISCUSSION

In this study, we isolated a B. thuringiensis DB27-resistant mutant in C. elegans that we showed to be identical to nasp-1, a gene that has previously been implicated in female development (51). Both nasp-1 alleles are reduction-of-function but not null alleles. Further characterization of *nasp-1* helped to discover a previously unknown function of *dcr-1*/Dicer in the immune response of *C*. elegans. Mammalian NASP1 is a tetratricopeptide (TPR) repeat domain-containing H1 linker histone binding protein that is part of a multichaperone complex implicated in nucleosome remodeling (64, 65). NASP1 is highly expressed in dividing cells and is required for normal cell cycle progression (65). A role for C. elegans nasp-1 in chromatin remodeling is supported by its interactions with a variety of chromatin-remodeling proteins like linker H1 histones and histone deacetylase (51). We hypothesize that nasp-1 might regulate the pathogen response of C. elegans via transcription and chromatin-remodeling mechanisms. Strikingly, it was shown that changes in NASP1 expression lead to the misregulation of a variety of genes in HeLa cells, including genes involved in the immune responses (66).

The RNAi deficiency of C. elegans nasp-1 is similar to that of other chromatin-remodeling genes (40) and provided the entry point for the discovery of the function of dcr-1 in C. elegans innate immunity, the most unexpected finding of this study. DCR-1 is an RNase III endoribonuclease whose primary function is the cleavage of dsRNA into smaller fragments that mediate RNAi (67). DCR-1 is required for miRNA processing and the synthesis of endo-siRNAs implicated in gene silencing and epigenetic regulation (67-69). Previous studies have shown a role for DCR-1 and the RNAi machinery in the C. elegans response to viral infections (14, 70). However, our results point toward a broader function of Dicer in innate immunity. To our knowledge, this study is the first to show a direct role for DCR-1 in antibacterial immunity. Results obtained by using different mutant alleles of C. elegans dcr-1 support the conclusion that its function in pathogen defense is mediated primarily by impaired miRNA biogenesis. Further evidence for this conclusion comes from the analysis of other genes that allow discrimination between miRNA processing and RNAi. Consistent with our conclusion is a recent report showing that miRNA activity in the C. elegans intestine is largely dedicated to attenuating the activity of pathogen response genes (48). Specifically, Kudlow et al. (48) have shown that ain-1 mutants, which have altered miRNA-induced silencing complex activity, are more resistant to P. aeruginosa than wild-type worms are. Additionally, those authors were able to identify specific miRNA families that are involved in damping of the immune response in the absence of infection. Together, these findings suggest that intestinal miRNAs act as negative regulators of infection response genes. However, we have been unable to identify a miRNA or siRNA that is likely to

be a part of the mechanisms of *nasp-1*-mediated resistance to *B*. *thuringiensis* DB27. Such studies might be complicated by the fact that redundancy among small RNAs is a well-known phenomenon in *C. elegans*.

The increased resistance of *dcr-1* mutants to *B. thuringiensis* DB27 might be due to the upregulation of pathogen response genes. Indeed, our RT-PCR experiment showed induction of some of the *B. thuringiensis* DB27-responsive genes in *dcr-1* mutants (Fig. 8A), indicating that DCR-1 may function as a suppressor of the C. elegans immune response to B. thuringiensis DB27 via suppression of miRNA synthesis. By using a reverse genetic approach, we found that one of the nasp-1-upregulated genes, namely, col-92, is an important determinant of the nasp-1 mutant's resistance that acts downstream of *nasp-1* and *dcr-1*. Collagens are differentially expressed in response to a variety of microbes (8, 71), suggesting that they may represent an essential part of the immune response. Nematodes, including C. elegans, are known for the unusually high number of collagens encoded in their genomes, and collagens have been implicated in life span and immunity (71–74). A potential role for collagens in the immune response might be in part responsible for the high number of nematode collagens, although direct proof of this assumption is hard to provide, even in the age of reverse genetics. One phenomenon that might hinder the identification of specific functions of collagens is redundancy, which is often seen among structurally related genes (73).

Our findings on the role of DCR-1 in C. elegans innate immunity strongly correlate with the identification of Dicer as a regulator of the mammalian immune response. Tang and coworkers found that the inhibition of Dicer in human cells upregulates major histocompatibility complex class I-related molecules A and B, which are innate immune system ligands for the NKG2D receptor expressed by natural killer cells (75). Those authors proposed further that a Dicer knockdown is indirectly linked to human innate immunity via the DNA damaging pathway. Whether a similar DNA-damaging pathway is involved in the upregulation of the C. elegans innate immune response awaits further analysis. Additionally, various miRNAs have been implicated in the regulation of diverse aspects of human innate immunity (76), including direct microbial killing. For example, miR-223-deficient neutrophils were shown to kill Candida albicans more effectively then wildtype cells do (77). A role for miRNAs in the human immune system is further supported by evidence that many immunological diseases are caused by alteration of miRNA function (76). Therefore, Dicer and miRNAs might play a conserved role as negative regulators in immunity.

Although alterations of *dcr-1* or miRNA functions are beneficial to *C. elegans* under pathogenic conditions because of immune response activation, this can be deleterious under normal growth conditions. Consistent with this, *nasp-1* mutants showed compromised reproduction, like many other pathogen-resistant mutants (47), indicating that the immune system might compete with the reproductive system for resources. Generally, chronic immune activation has been linked to several pathological disorders, like inflammatory bowel disease in humans (78). Therefore, it is crucial to any organism to have mechanisms of innate immune suppression in the absence of pathogens.

On the basis of our results, we propose the following mechanism for the function of *nasp-1* in the regulation of *C. elegans* innate immunity (Fig. 8F). First, *nasp-1* is a regulator of *dcr-1*. In

nasp-1 mutants, dcr-1 function is repressed because of chromatin remodeling or related mechanisms, which leads to, besides RNAi deficiency, the inhibition of dcr-1-mediated miRNA processing. Consequently, certain miRNAs are not processed and, in turn, target genes, including col-92, are more strongly expressed than normal. As a consequence, mutant worms are more resistant to B. thuringiensis DB27 infection. Additionally, nasp-1, dcr-1, and drsh-1 mutants exhibit increased resistance to S. aureus as a secondary consequence of altered reproduction likely because of germ line defects. In general, germ line defects and fecundity might play a role in the observed resistance phenotypes of nasp-1 and dcr-1 mutants. It is important to note, however, that nasp-1 dcr-1 double mutants with cdc-25 RNAi gave different results for resistance to B. thuringiensis DB27 and S. aureus, respectively. These findings indicate complex interactions between fecundity and nematode immunity.

Given the comparative framework of our work, what about the specificity of nasp-1 and the resistance of P. pacificus to B. thuringiensis DB27? While *nasp-1* is clearly conserved in sequence, the amino acid positions that are changed in the two C. elegans nasp-1 mutants are not. It is therefore possible that the exact molecular mechanisms of nasp-1 and its role in nematode immunity did change during the course of evolution. More generally, our parallel work on innate immunity in P. pacificus has recently revealed that more general anatomical and physiological parameters are in part responsible for the observed differences between the two species. P. pacificus is not equipped with a grinder, as is typical of members of the family Diplogastridae, and so does not lyse its bacterial food in the pharynx. P. pacificus mutants hypersusceptible to bacterial pathogens are often Unc mutants with an abnormal defecation cycle, indicating that the regulation of the defecation cycle is crucial for innate immunity (32). Together, this comparative work on innate nematode immunity points toward the correlated action of anatomical, physiological, and molecular mechanisms that control nematode-bacterial interactions and regulate the evolution of innate nematode immunity.

ACKNOWLEDGMENTS

We are grateful to Gabi Eberhardt for help with the sequencing of *btr-1* mutant. We thank Adrian Streit and Robbie Rae and for critical reading of the manuscript and invaluable discussions and feedback.

We thank the *Caenorhabditis* Genetics Center, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440), for providing strains.

This research was supported by the Max Planck Society and by the DFG-funded RTG 1708, "Molecular principles of bacterial survival strategies."

REFERENCES

- 1. Irazoqui JE, Urbach JM, Ausubel FM. 2010. Evolution of host innate defence: insights from Caenorhabditis elegans and primitive invertebrates. Nat. Rev. Immunol. 10:47–58.
- Pukkila-Worley R, Ausubel FM. 2012. Immune defense mechanisms in the Caenorhabditis elegans intestinal epithelium. Curr. Opin. Immunol. 24:3–9.
- Tan M-W, Shapira M. 2011. Genetic and molecular analysis of nematode-microbe interactions. Cell. Microbiol. 13:497–507.
- 4. Partridge FA, Gravato-Nobre MJ, Hodgkin J. 2010. Signal transduction pathways that function in both development and innate immunity. Dev. Dyn. 239:1330–1336.
- Schulenburg H, Kurz CL, Ewbank JJ. 2004. Evolution of the innate immune system: the worm perspective. Immunol. Rev. 198:36–58.
- Engelmann I, Pujol N. 2010. Innate immunity in C. elegans. Adv. Exp. Med. Biol. 708:105–121.

- Kim DH, Ausubel FM. 2005. Evolutionary perspectives on innate immunity from the study of Caenorhabditis elegans. Curr. Opin. Immunol. 17:4–10.
- Wong D, Bazopoulou D, Pujol N, Tavernarakis N, Ewbank J. 2007. Genome-wide investigation reveals pathogen-specific and shared signatures in the response of Caenorhabditis elegans to infection. Genome Biol. 8:R194. doi:10.1186/gb-2007-8-9-r194.
- O'Rourke D, Baban D, Demidova M, Mott R, Hodgkin J. 2006. Genomic clusters, putative pathogen recognition molecules, and antimicrobial genes are induced by infection of C. elegans with M. nematophilum. Genome Res. 16:1005–1016.
- Engelmann I, Griffon A, Tichit L, Montañana-Sanchis F, Wang G, Reinke V, Waterston RH, Hillier LW, Ewbank JJ. 2011. A comprehensive analysis of gene expression changes provoked by bacterial and fungal infection in C. elegans. PLoS One 6:e19055. doi:10.1371/journal.pone .0019055.
- Sinha A, Rae R, Iatsenko I, Sommer RJ. 2012. System wide analysis of the evolution of innate immunity in the nematode model species Caenorhabditis elegans and Pristionchus pacificus. PLoS One 7:e44255. doi:10.1371 /journal.pone.0044255.
- Marsh EK, May RC. 2012. *Caenorhabditis elegans*, a model organism for investigating immunity. Appl. Environ. Microbiol. 78:2075–2081.
- Pujol N, Zugasti O, Wong D, Couillault C, Kurz CL, Schulenburg H, Ewbank JJ. 2008. Anti-fungal innate immunity in C. elegans is enhanced by evolutionary diversification of antimicrobial peptides. PLoS Pathog. 4:e1000105. doi:10.1371/journal.ppat.1000105.
- 14. Félix M-A, Ashe A, Piffaretti J, Wu G, Nuez I, Bélicard T, Jiang Y, Zhao G, Franz CJ, Goldstein LD, Sanroman M, Miska EA, Wang D. 2011. Natural and experimental infection of Caenorhabditis nematodes by novel viruses related to nodaviruses. PLoS Biol. 9:e1000586. doi:10.1371 /journal.pbio.1000586.
- Troemel ER. 2011. New models of microsporidiosis: infections in zebrafish, C. elegans, and honey bee. PLoS Pathog. 7:e1001243. doi:10.1371 /journal.ppat.1001243.
- Feinbaum RL, Urbach JM, Liberati NT, Djonovic S, Adonizio A, Carvunis A-R, Ausubel FM. 2012. Genome-wide identification of Pseudomonas aeruginosa virulence-related genes using a Caenorhabditis elegans infection model. PLoS Pathog. 8:e1002813. doi:10.1371/journal .ppat.1002813.
- Kurz CL, Chauvet S, Andrès E, Aurouze M, Vallet I, Michel GPF, Uh M, Celli J, Filloux A, De Bentzmann S, Steinmetz I, Hoffmann JA, Finlay BB, Gorvel J-P, Ferrandon D, Ewbank JJ. 2003. Virulence factors of the human opportunistic pathogen Serratia marcescens identified by in vivo screening. EMBO J. 22:1451–1460.
- Schulte RD, Makus C, Hasert B, Michiels NK, Schulenburg H. 2010. Multiple reciprocal adaptations and rapid genetic change upon experimental coevolution of an animal host and its microbial parasite. Proc. Natl. Acad. Sci. U. S. A. 107:7359–7364.
- Bravo A, Gill SS, Soberón M. 2007. Mode of action of Bacillus thuringiensis Cry and Cyt toxins and their potential for insect control. Toxicon 49:423–435.
- Wei J-Z, Hale K, Carta L, Platzer E, Wong C, Fang S-C, Aroian RV. 2003. Bacillus thuringiensis crystal proteins that target nematodes. Proc. Natl. Acad. Sci. U. S. A. 100:2760–2765.
- Huffman DL, Bischof LJ, Griffitts JS, Aroian RV. 2004. Pore worms: using Caenorhabditis elegans to study how bacterial toxins interact with their target host. Int. J. Med. Microbiol. 293:599–607.
- Griffitts JS, Whitacre JL, Stevens DE, Aroian RV. 2001. Bt toxin resistance from loss of a putative carbohydrate-modifying enzyme. Science 293:860–864.
- Huffman DL, Abrami L, Sasik R, Corbeil J, Van der Goot FG, Aroian RV. 2004. Mitogen-activated protein kinase pathways defend against bacterial pore-forming toxins. Proc. Natl. Acad. Sci. U. S. A. 101:10995– 11000.
- Bellier A, Chen C-S, Kao C-Y, Cinar HN, Aroian RV. 2009. Hypoxia and the hypoxic response pathway protect against pore-forming toxins in C. elegans. PLoS Pathog. 5:e1000689. doi:10.1371/journal.ppat.1000689.
- Bischof LJ, Kao C-Y, Los FCO, Gonzalez MR, Shen Z, Briggs SP, Van der Goot FG, Aroian RV. 2008. Activation of the unfolded protein response is required for defenses against bacterial pore-forming toxin in vivo. PLoS Pathog. 4:e1000176. doi:10.1371/journal.ppat.1000176.
- 26. Kao C-Y, Los FCO, Huffman DL, Wachi S, Kloft N, Husmann M, Karabrahimi V, Schwartz J-L, Bellier A, Ha C, Sagong Y, Fan H, Ghosh

P, Hsieh M, Hsu C-S, Chen L, Aroian RV. 2011. Global functional analyses of cellular responses to pore-forming toxins. PLoS Pathog. 7:e1001314. doi:10.1371/journal.ppat.1001314.

- 27. Los FCO, Kao C-Y, Smitham J, McDonald KL, Ha C, Peixoto CA, Aroian RV. 2011. RAB-5- and RAB-11-dependent vesicle-trafficking pathways are required for plasma membrane repair after attack by bacterial pore-forming toxin. Cell Host Microbe 9:147–157.
- Félix M-A, Duveau F. 2012. Population dynamics and habitat sharing of natural populations of Caenorhabditis elegans and C. briggsae. BMC Biol. 10:59. doi:10.1186/1741-7007-10-59.
- Herrmann M, Kienle S, Rochat J, Mayer WE, Sommer RJ. 2010. Haplotype diversity of the nematode Pristionchus pacificus on Réunion in the Indian Ocean suggests multiple independent invasions. Biol. J. Linn. Soc. 100:170–179.
- Rae R, Riebesell M, Dinkelacker I, Wang Q, Herrmann M, Weller AM, Dieterich C, Sommer RJ. 2008. Isolation of naturally associated bacteria of necromenic Pristionchus nematodes and fitness consequences. J. Exp. Biol. 211:1927–1936.
- Rae R, Iatsenko I, Witte H, Sommer RJ. 2010. A subset of naturally isolated Bacillus strains show extreme virulence to the free-living nematodes Caenorhabditis elegans and Pristionchus pacificus. Environ. Microbiol. 12:3007–3021.
- Rae R, Witte H, Rödelsperger C, Sommer RJ. 2012. The importance of being regular: Caenorhabditis elegans and Pristionchus pacificus defecation mutants are hypersusceptibile to bacterial pathogens. Int. J. Parasitol. 42:747–753.
- 33. Garsin DA, Villanueva JM, Begun J, Kim DH, Sifri CD, Calderwood SB, Ruvkun G, Ausubel FM. 2003. Long-lived C. elegans daf-2 mutants are resistant to bacterial pathogens. Science 300:1921. doi:10.1126/science .1080147.
- Fay D. 14 June 2006, posting date. Genetic mapping and manipulation: chapter 7—making compound mutants. *In* The *C. elegans* Research Community (ed), WormBook, http://www.wormbook.org. doi/10.1895/wormbook.1.96.2.
- 35. Davis MW, Hammarlund M, Harrach T, Hullett P, Olsen S, Jorgensen EM. 2005. Rapid single nucleotide polymorphism mapping in C. elegans. BMC Genomics 6:118. doi:10.1186/1471-2164-6-118.
- Mello CC, Kramer JM, Stinchcomb D, Ambros V. 1991. Efficient gene transfer in C. elegans: extrachromosomal maintenance and integration of transforming sequences. EMBO J. 10:3959–3970.
- Hobert O. 2002. PCR fusion-based approach to create reporter gene constructs for expression analysis in transgenic C. elegans. Biotechniques 32: 728–730.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754–1760.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25:2078–2079.
- 40. Cui M, Kim EB, Han M. 2006. Diverse chromatin remodeling genes antagonize the Rb-involved SynMuv pathways in C. elegans. PLoS Genet. 2:e74. doi:10.1371/journal.pgen.0020074.
- Sinha A, Sommer RJ, Dieterich C. 2012. Divergent gene expression in the conserved dauer stage of the nematodes Pristionchus pacificus and Caenorhabditis elegans. BMC Genomics 13:254. doi:10.1186/1471-2164-13 -254.
- 42. Harris TW, Antoshechkin I, Bieri T, Blasiar D, Chan J, Chen WJ, De La Cruz N, Davis P, Duesbury M, Fang R, Fernandes J, Han M, Kishore R, Lee R, MüllerH-M, Nakamura C, Ozersky P, Petcherski A, Rangarajan A, Rogers A, Schindelman G, Schwarz EM, Tuli MA, Van Auken K, Wang D, Wang X, Williams G, Yook K, Durbin R, Stein LD, Spieth J, Sternberg PW. 2010. WormBase: a comprehensive resource for nematode research. Nucleic Acids Res. 38:D463–D467.
- Powell JR, Kim DH, Ausubel FM. 2009. The G protein-coupled receptor FSHR-1 is required for the Caenorhabditis elegans innate immune response. Proc. Natl. Acad. Sci. U. S. A. 106:2782–2787.
- 44. Shivers RP, Pagano DJ, Kooistra T, Richardson CE, Reddy KC, Whitney JK, Kamanzi O, Matsumoto K, Hisamoto N, Kim DH. 2010. Phosphorylation of the conserved transcription factor ATF-7 by PMK-1 p38 MAPK regulates innate immunity in Caenorhabditis elegans. PLoS Genet. 6:e1000892. doi:10.1371/journal.pgen.1000892.
- Richardson CE, Kooistra T, Kim DH. 2010. An essential role for XBP-1 in host protection against immune activation in C. elegans. Nature 463: 1092–1095.
- 46. Fuhrman LE, Goel AK, Smith J, Shianna KV, Aballay A. 2009. Nucleolar

proteins suppress Caenorhabditis elegans innate immunity by inhibiting p53/CEP-1. PLoS Genet. 5:e1000657. doi:10.1371/journal.pgen.1000657.

- Miyata S, Begun J, Troemel ER, Ausubel FM. 2008. DAF-16-dependent suppression of immunity during reproduction in Caenorhabditis elegans. Genetics 178:903–918.
- Kudlow BA, Zhang L, Han M. 2012. Systematic analysis of tissuerestricted miRISCs reveals a broad role for microRNAs in suppressing basal activity of the C. elegans pathogen response. Mol. Cell 46:530–541.
- Ward S, Miwa J. 1978. Characterization of temperature-sensitive, fertilization-defective mutants of the nematode caenorhabditis elegans. Genetics 88:285–303.
- Sarin S, Prabhu S, O'Meara MM, Pe'er I, Hobert O. 2008. Caenorhabditis elegans mutant allele identification by whole-genome sequencing. Nat. Methods 5:865–867.
- Grote P, Conradt B. 2006. The PLZF-like protein TRA-4 cooperates with the Gli-like transcription factor TRA-1 to promote female development in C. elegans. Dev. Cell 11:561–573.
- 52. Kim DH, Feinbaum R, Alloing G, Emerson FE, Garsin DA, Inoue H, Tanaka-Hino M, Hisamoto N, Matsumoto K, Tan M-W, Ausubel FM. 2002. A conserved p38 MAP kinase pathway in Caenorhabditis elegans innate immunity. Science 297:623–626.
- 53. Mizuno T, Hisamoto N, Terada T, Kondo T, Adachi M, Nishida E, Kim DH, Ausubel FM, Matsumoto K. 2004. The Caenorhabditis elegans MAPK phosphatase VHP-1 mediates a novel JNK-like signaling pathway in stress response. EMBO J. 23:2226–2234.
- Tijsterman M, May RC, Simmer F, Okihara KL, Plasterk RHA. 2004. Genes required for systemic RNA interference in Caenorhabditis elegans. Curr. Biol. 14:111–116.
- 55. Kim JK, Gabel HW, Kamath RS, Tewari M, Pasquinelli A, Rual J-F, Kennedy S, Dybbs M, Bertin N, Kaplan JM, Vidal M, Ruvkun G. 2005. Functional genomic analysis of RNA interference in C. elegans. Science 308:1164–1167.
- Tabara H, Sarkissian M, Kelly WG, Fleenor J, Grishok A, Timmons L, Fire A, Mello CC. 1999. The rde-1 gene, RNA interference, and transposon silencing in C. elegans. Cell 99:123–132.
- Welker NC, Habig JW, Bass BL. 2007. Genes misregulated in C. elegans deficient in Dicer, RDE-4, or RDE-1 are enriched for innate immunity genes. RNA 13:1090–1102.
- Mori MA, Raghavan P, Thomou T, Boucher J, Robida-Stubbs S, Macotela Y, Russell SJ, Kirkland JL, Blackwell TK, Kahn CR. 2012. Role of microRNA processing in adipose tissue in stress defense and longevity. Cell Metab. 16:336–347.
- TeKippe M, Aballay A. 2010. C. elegans germline-deficient mutants respond to pathogen infection using shared and distinct mechanisms. PLoS One 5:e11777. doi:10.1371/journal.pone.0011777.
- Arantes-Oliveira N, Apfeld J, Dillin A, Kenyon C. 2002. Regulation of life-span by germ-line stem cells in Caenorhabditis elegans. Science 295: 502–505.
- Pavelec DM, Lachowiec J, Duchaine TF, Smith HE, Kennedy S. 2009. Requirement for the ERI/DICER complex in endogenous RNA interference and sperm development in Caenorhabditis elegans. Genetics 183: 1283–1295.
- 62. Bouasker S, Simard MJ. 2012. The slicing activity of miRNA-specific Argonautes is essential for the miRNA pathway in C. elegans. Nucleic Acids Res. 40:10452–10462.
- Denli AM, Tops BBJ, Plasterk RHA, Ketting RF, Hannon GJ. 2004. Processing of primary microRNAs by the Microprocessor complex. Nature 432:231–235.
- Richardson R, Batova IN, Widgren EE, Zheng LX, Whitfield M, Marzluff WF, O'Rand MG. 2000. Characterization of the histone H1-binding protein, NASP, as a cell cycle-regulated somatic protein. J. Biol. Chem. 275:30378–30386.
- 65. Richardson RT, Alekseev OM, Grossman G, Widgren EE, Thresher R, Wagner EJ, Sullivan KD, Marzluff WF, O'Rand MG. 2006. Nuclear autoantigenic sperm protein (NASP), a linker histone chaperone that is required for cell proliferation. J. Biol. Chem. 281:21526–21534.
- Alekseev OM, Richardson RT, Alekseev O, O'Rand MG. 2009. Analysis of gene expression profiles in HeLa cells in response to overexpression or siRNA-mediated depletion of NASP. Reprod. Biol. Endocrinol. 7:45. doi: 10.1186/1477-7827-7-45.
- Jinek M, Doudna JA. 2009. A three-dimensional view of the molecular machinery of RNA interference. Nature 457:405–412.

- Guo H, Ingolia NT, Weissman JS, Bartel DP. 2010. Mammalian microRNAs predominantly act to decrease target mRNA levels. Nature 466:835–840.
- 69. Duchaine TF, Wohlschlegel JA, Kennedy S, Bei Y, Conte D, Jr, Pang K, Brownell DR, Harding S, Mitani S, Ruvkun G, Yates JR, III, Mello CC. 2006. Functional proteomics reveals the biochemical niche of C. elegans DCR-1 in multiple small-RNA-mediated pathways. Cell 124: 343–354.
- Lu R, Maduro M, Li F, Li HW, Broitman-Maduro G, Li WX, Ding SW. 2005. Animal virus replication and RNAi-mediated antiviral silencing in Caenorhabditis elegans. Nature 436:1040–1043.
- Coolon JD, Jones KL, Todd TC, Carr BC, Herman MA. 2009. Caenorhabditis elegans genomic response to soil bacteria predicts environmentspecific genetic effects on life history traits. PLoS Genet. 5:e1000503. doi: 10.1371/journal.pgen.1000503.
- Kenning C, Kipping I, Sommer RJ. 2004. Isolation of mutations with dumpy-like phenotypes and of collagen genes in the nematode Pristionchus pacificus. Genesis 40:176–183.
- 73. Page AP, Johnstone IL. 19 March 2007, posting date. The cuticle. In

The *C. elegans* Research Community (ed), WormBook, http://www .wormbook.org. doi/10.1895/wormbook.1.138.1.

- 74. Sahu SN, Lewis J, Patel I, Bozdag S, Lee JH, LeClerc JE, Cinar HN. 2012. Genomic analysis of immune response against Vibrio cholerae hemolysin in Caenorhabditis elegans. PLoS One 7:e38200. doi:10.1371/journal.pone .0038200.
- Tang K-F, Ren H, Cao J, Zeng G-L, Xie J, Chen M, Wang L, He C-X. 2008. Decreased Dicer expression elicits DNA damage and up-regulation of MICA and MICB. J. Cell Biol. 182:233–239.
- O'Connell RM, Rao DS, Chaudhuri AA, Baltimore D. 2010. Physiological and pathological roles for microRNAs in the immune system. Nat. Rev. Immunol. 10:111–122.
- Johnnidis JB, Harris MH, Wheeler RT, Stehling-Sun S, Lam MH, Kirak O, Brummelkamp TR, Fleming MD, Camargo FD. 2008. Regulation of progenitor cell proliferation and granulocyte function by microRNA-223. Nature 451:1125–1129.
- Cario E. 2010. Heads up! How the intestinal epithelium safeguards mucosal barrier immunity through the inflammasome and beyond. Curr. Opin. Gastroenterol. 26:583–590.